

6-22-2017

New insight into the cluster structure of ^9Be by reactions with deuteron beam

S. M. Lukyanov

Flerov Laboratory of Nuclear Reactions, Russian Federation

A. S. Denikin

Flerov Laboratory of Nuclear Reactions, Russian Federation

M. A. Naumenko

Flerov Laboratory of Nuclear Reactions, Russian Federation

V. Burjan

Nuclear Physics Institute, Czech Republic

W. H. Trzaska

Department of Physics, University of Jyväskylä, Finland

See next page for additional authors

Follow this and additional works at: <https://www.ephys.kz/journal>



Part of the [Materials Science and Engineering Commons](#), and the [Physics Commons](#)

Recommended Citation

Lukyanov, S. M.; Denikin, A. S.; Naumenko, M. A.; Burjan, V.; Trzaska, W. H.; Harakeh, M.; Etasse, D.; Kroha, V.; Mrazek, J.; Mendibayev, K.; Sivacek, I.; Glagolev, V.; Piskor, S.; Penionzhkevich, Yu. E.; Skobelev, N. K.; Stefan, I.; Verney, D.; Kuterbekov, K. A.; and Zholdybayev, T. (2017) "New insight into the cluster structure of ^9Be by reactions with deuteron beam," *Eurasian Journal of Physics and Functional Materials*: Vol. 1: No. 2, Article 4.

DOI: <https://doi.org/10.29317/ejpfm.2017010204>

This Original Study is brought to you for free and open access by Eurasian Journal of Physics and Functional Materials. It has been accepted for inclusion in Eurasian Journal of Physics and Functional Materials by an authorized editor of Eurasian Journal of Physics and Functional Materials.

New insight into the cluster structure of ^9Be by reactions with deuteron beam

Authors

S. M. Lukyanov, A. S. Denikin, M. A. Naumenko, V. Burjan, W. H. Trzaska, M. Harakeh, D. Etasse, V. Kroha, J. Mrazek, K. Mendibayev, I. Sivacek, V. Glagolev, S. Piskor, Yu. E. Penionzhkevich, N. K. Skobelev, I. Stefan, D. Verney, K. A. Kuterbekov, and T. Zholdybayev

New insight into the cluster structure of ${}^9\text{Be}$ by reactions with deuteron beam

S.M. Lukyanov¹, A.S. Denikin¹, M.A. Naumenko¹, V. Burjan²,
W.H. Trzaska³, M. Harakeh⁶, D. Etasse⁷, V. Kroha², J. Mrazek²,
K. Mendibayev^{1,4,5}, I. Sivaček², V. Glagolev², Š. Piskoř²,
Yu.E. Penionzhkevich^{1,9}, N.K. Skobelev¹, I. Stefan⁸, D. Verney⁸,
K.A. Kuterbekov⁴, T. Zholdybayev⁴

¹Flerov Laboratory of Nuclear Reactions, Dubna, Russian Federation

²Nuclear Physics Institute, Řež, Czech Republic

³Department of Physics, University of Jyväskylä, Jyväskylä, Finland

⁴L.N. Gumilyov Eurasian National University, Astana, Kazakhstan

⁵Institute of Nuclear Physics, Almaty, Kazakhstan

⁶KVI-CART, University of Groningen, Groningen, The Netherlands

⁷LPC-Caen, ENSICAEN, Université de Caen, CNRS/IN2P3-ENSI, Caen, France

⁸Institut de Physique Nucléaire, Univ. Paris-Sud, Université Paris-Saclay, Orsay, France

⁹National Research Nuclear University "MEPhI", Moscow, Russian Federation

*e-mail: lukyan@jinr.ru

DOI: 10.29317/ejpfm.2017010204

Received: 17.11.2017

Angular distributions of protons, deuterons, tritons and alpha particles emitted in the reaction ${}^2\text{H} + {}^9\text{Be}$ at $E_{lab} = 19.5$ MeV were measured with an aim to shed light on the internal cluster structure of ${}^9\text{Be}$ and to study possible cluster transfer of ${}^5\text{He}$. The analyses suggest a significant contribution of five-nucleon transfer in the reaction channel ${}^9\text{Be}(d, {}^4\text{He}){}^7\text{Li}$.

Keywords: cluster structure, transfer in the reaction channel.

Introduction

Due to the Borromean structure, a special attention has been focused on the ${}^9\text{Be}$ nucleus, the breakup of which can occur directly into two alpha particles and a neutron or via one of two unstable intermediate nuclei such as ${}^8\text{Be}$ or ${}^5\text{He}$ [1,2]. Scattering of a projectile, such as ${}^1,2\text{H}$ or ${}^3,4\text{He}$, on a target is a standard tool to study the structure of nuclei. This method involves the angular distribution measurement of elastic and inelastic scattering. Energy distribution of the products gives information about internal structure of the interacting nuclei. The angular distributions of the ${}^9\text{Be}({}^3\text{He}, {}^3\text{He}){}^9\text{Be}$, ${}^9\text{Be}({}^3\text{He}, {}^5\text{He}){}^7\text{Be}$, ${}^9\text{Be}({}^3\text{He}, {}^5\text{Li}){}^7\text{Li}$, ${}^9\text{Be}({}^3\text{He}, {}^6\text{Be}){}^6\text{He}$, and ${}^9\text{Be}({}^3\text{He}, {}^6\text{Li}){}^6\text{Li}$ reaction channels were measured in [3,4] and described within the framework of the optical model, the coupled-channel approach and the distorted-wave Born approximation. The performed analysis of the experimental data shows that the potential parameters are quite sensitive to the exit channel and hence to the cluster structure of the populated

states, which allows one to make general observations and conclusions regarding the internal structure of the target and product nuclei. The experiment [3,4] was designed to study the breakup of ${}^9\text{Be}$ in an attempt to determine the contribution of the ${}^8\text{Be}+n$ and ${}^5\text{He}+\alpha$ channels in the inclusive measurements. We found that the ratio about 2.7:1 may be assigned to the contributions of these two channels, respectively. The determined value justifies that the ${}^5\text{He}+\alpha$ breakup channel plays an important role.

Another aspect is an attempt to find not only the cluster structure (for instance, ${}^5\text{He}$), but also clarify how the cluster structure is involved into nuclear reaction mechanism. Indeed, starting from Detraz [5,6] multiparticle-multihole structures were expected to occur at rather low excitation energies in nuclei. Four-nucleon transfer reactions are being extensively studied. One may hope that their major features, in spite of the a-priori complexity of such a transfer, can be understood assuming that the nucleons are transferred as a whole, strongly correlated in a cluster which has the internal quantum numbers of a free particle.

Experimental Method

The experiment was performed using 19.5 MeV beam energy of ${}^2\text{H}$ ions on the cyclotron of the INP (Řež, Czech Republic). The average beam current during the experiment was maintained at 10 nA. The self-supporting Be target was prepared from a thin beryllium foil of 99 % purity. To measure (in)elastically scattered ions, a set of four telescopes each consisting of ΔE_0 , ΔE , E_r detectors with thicknesses of 12, 100 μm and 3 mm, respectively, were used. The telescopes were mounted at a distance of about 19 cm from the target in a reaction chamber. Particle identification was performed based on the energy-loss measurements of ΔE and residual energy E_r , i.e., by the so-called $\Delta E \times E$ method. An example of two-dimensional plots (yield vs. energy loss ΔE and residual energy E_r) is shown in Figure 1.




Figure 1. Particle identification plots for the products of the ${}^2\text{H}+{}^9\text{Be}$ reaction: p , d , t , and ${}^4\text{He}$. ΔE is the energy loss and E_r is the residual energy. Excited states for the ${}^7\text{Li}+\alpha$ reaction channel ${}^7\text{Li}+\alpha$ are indicated.

This experimental technique allows us to identify the particles p , d , t , and ${}^4\text{He}$ and determine their total deposited energies. The spectra of total deposited energy are shown in Figure 2. All peaks, which can be observed in the histograms in

Figure 2, were identified and found to belong to the ground and the excited states of ^{10}Be , ^9Be , ^8Be , and ^7Li , as the complementary products to the detected particles p , d , t , and ^4He , respectively.




Figure 2. Total deposited energy spectra measured at $\theta_{lab} = 32^\circ$ for the detected p (panel a), d (panel b), t (panel c), and ^4He (panel d). The ground and the excited states of ^7Li for the case of detected complementary product ^4He as well as the ground states and the excited states for ^8Be , ^9Be , and ^{10}Be in the case of detected t , d and p as complementary products, respectively, were unambiguously identified.

Results and Data Analysis

The differential cross section for the elastic, inelastic, and transfer reaction channels are presented in Figure 3. All calculations and fitting of the experimental data have been performed using the FRESKO [7] code in the framework of the CC method.

The calculation for (in)elastic scattering (Figure 3a) was performed using the method of strong channel coupling in the adiabatic rotational model. It is assumed that the state 2.43 MeV is the first excited state in the rotational band of ^9Be . The quadrupole deformation parameter of the target nucleus found from the performed analysis of data is $\beta_2 = 0.64$, which perfectly coincides with the previous analysis of inelastic scattering of ^4He nuclei on the same target [3]. The optical potential parameters used for the calculation are similar to those obtained in [8]. Good agreement between experimental data and calculations is observed, as shown in Figure 3a.

A special attention was paid to the $^4\text{He} + ^7\text{Li}$ reaction channel, which can proceed through two different mechanisms: either transfer of d (dashed line in Figure 3b) or transfer of ^5He (solid line in Figure 3b) from the target to the projectile. In the exit channel both these mechanisms are indistinguishable from each other. However, in the former case the ^4He nuclei are expected to fly in the forward direction in the center of mass system, whereas in the latter case the alpha particles




Figure 3. a) Experimental angular distribution for elastic (\square) and inelastic scattering (\circ) of 19.5 MeV deuterons from ^9Be target compared with the calculations. The curves are the results of the optical-model and the coupled-channel calculations. b) Experimental angular distribution (\bullet) for $^4\text{He} + ^7\text{Li}$ reaction channel (see details in text).

should preferably fly at the backward angles in the center of mass system. The calculations carried out within the DWBA method for these two mechanisms are shown as the dashed and solid curves in Figure 3b. Their coherent sum shown as the black curve is in good agreement with the experimental points. It is interesting to note that for the correct description of the data, it is necessary to set the sufficiently large values of the spectroscopic factors for the systems $^9\text{Be} = \alpha + ^5\text{He}$ ($S=1.2$) and $^7\text{Li} = d + ^5\text{He}$ ($S=1.0$). In addition, to describe the structure of the angular distributions it is also necessary to assume a 30% admixture of the d -state in the structure of ^7Li .

In summary, angular distributions for the $^9\text{Be}(d, d)^9\text{Be}^*$, $^9\text{Be}(d, p)^{10}\text{Be}$, $^9\text{Be}(d, t)^8\text{Be}$, and $^9\text{Be}(d, ^4\text{He})^7\text{Li}$ channels were measured. Experimental angular distributions were described within the optical model, the coupled channel approach, and the distorted wave Born approximation. The optical model provides good agreement with the elastic scattering. The DWBA calculations agree well with the transfer reaction data. The spectroscopic factors for the systems $^9\text{Be} = \alpha + ^5\text{He}$ and $^7\text{Li} = d + ^5\text{He}$ are close to unity, which confirms significant contribution of the considered cluster configurations to the structure of ground states. The analysis shows that the contribution of the compound nucleus mechanism is negligible. In the $(d, ^4\text{He})$ channel (see Figure 3), the deuteron transfer provides only a small contribution, whereas a relatively large contribution of ^5He transfer was found in agreement with the result [8]. This demonstrates that the specific structure of the ^9Be nucleus as a weakly bound system of two alpha particles and a neutron strongly favors the five-nucleon transfer compared to the deuteron transfer.

Acknowledgments

The authors thank the CANAM project (<http://users.canam.ujf.cas.cz>) for providing beam time for the experiment and support. This work was supported by Russian Science Foundation (17-12-01170).

References

- [1] T.A.D. Brown et al., Phys. Rev. C **76**, (2007) 054605.
- [2] P. Papka et al., Phys. Rev. C **75**, (2007) 045803.
- [3] S.M. Lukyanov et al., J. Phys. G **41** (2014) 035103.
- [4] S.M. Lukyanov et al., World Journal of Nuclear Science and Technology **5** (2015) 265.
- [5] C. Detraz et al., Nucl. Phys. A **147** (1970) 488.
- [6] C. Detraz et al., Nucl. Phys. A **228** (1974) 39.
- [7] <http://www.ianthompson.org/surrey> and <http://nrv.jinr.ru>.
- [8] A. Szczurek et al., Zeitschrift fur Physik A Atomic Nuclei, **333** (1989) 271.